

\$\$\$STN:Highlighton= ***Highlightoff=***
Connecting via Winsock to STN

Welcome to STN International! Enter x:x

LOGINID:SSSPTA1653RAW

PASSWORD:

TERMINAL (ENTER 1, 2, 3, OR ?):2

***** Welcome to STN International *****

NEWS 1 Web Page URLs for STN Seminar Schedule - N. America
NEWS 2 "Ask CAS" for self-help around the clock
NEWS 3 JUL 12 BRILSTEIN enhanced with new display and select options,
resulting in a closer connection to BABS
NEWS 4 AUG 02 IFIPAT/IFIUDB/IFICDB reloaded with new search and display
fields
NEWS 5 AUG 02 Caplus and CA patent records enhanced with European and Japan
Patent Office Classifications
NEWS 6 AUG 02 The Analysis Edition of STN Express with Discover!
(version 7.01 for Windows) now available
NEWS 7 AUG 27 BIOCOMMERCE: Changes and enhancements to content coverage
NEWS 8 AUG 27 BIOTECCHAS/BIOTECCHDS: Two new display fields added for legal
status data from INPADOC
NEWS 9 SEP 01 INPADOC: New family current-awareness alert (SDI) available
NEWS 10 SEP 01 New pricing for the Save Answers for SciFinder Wizard within
STN Express with Discover!
NEWS 11 SEP 01 New display format, HITSTR, available in WPIDS/WPINDEX/WPIX
NEWS 12 SEP 27 STANDARDS will no longer be available on STN
NEWS 13 SEP 27 SWETSCAN will no longer be available on STN
NEWS 14 OCT 28 KOREAPAT now available on STN
NEWS EXPRESS OCTOBER 29 CURRENT WINDOWS VERSION IS V7.01A, CURRENT
MACINTOSH VERSION IS V6.0C(ENG) AND V6.0JC(JP),
AND CURRENT DISCOVER FILE IS DATED 11 AUGUST 2004
NEWS HOURS STN Operating Hours Plus Help Desk Availability
NEWS INTER General Internet Information
NEWS LOGIN Welcome Banner and News Items
NEWS PHONE Direct Dial and Telecommunication Network Access to STN
NEWS WWW CAS World Wide Web Site (general information)

Enter NEWS followed by the item number or name to see news on that
specific topic.

All use of STN is subject to the provisions of the STN Customer
agreement. Please note that this agreement limits use to scientific
research. Use for software development or design or implementation
of commercial gateways or other similar uses is prohibited and may
result in loss of user privileges and other penalties.

***** STN Columbus *****

FILE 'HOME' ENTERED AT 18:48:02 ON 07 NOV 2004

=> index biosci

FILE 'DRUGMONOC' ACCESS NOT AUTHORIZED

COST IN U.S. DOLLARS

FULL ESTIMATED COST

INDEX	ADISCTI, ADISINSIGHT, ADISNEWS, AGRICOLA, ANABSTR, ANTE, AQUALINE,
AQUASCI, BIOBUSINESS, BIOCOMMERCE, BIOENG, BIOSIS, BIOTECCHAS, BIOTECCHDS,	
BIOTECCHNO, CAB, CANCERLIT, CAPLUS, CEBA-VTB, CEN, CIN, CONFSCI, CROPB,	
CROPU, DDBF, DDFU, DGENE, DISSABS, ...	

2004

75 FILES IN THE FILE LIST IN STNINDEX

Enter SET DETAIL ON to see search term postings or to view
search error messages that display as 0* with SET DETAIL OFF.

=> s (bacteriophage or phage) (w) 77

1	FILE ADISINSIGHT
1	FILE BIOENG
10	FILE BIOSIS
2	FILE BIOTECCHAS
2	FILE BIOTECCHDS
1	FILE BIOTECCHNO
11	FILE CAPLUS

26 FILES SEARCHED...

25	FILE DGENE
2	FILE DISSABS
4	FILE EMBASE
2	FILE EMBIOMBASE
24	FILE GENBANK
3	FILE IFIPAT
2	FILE LIFESCI
4	FILE MEDLINE
2	FILE PASCAL
1	FILE PHAR

58 FILES SEARCHED...

2	FILE PROMT
3	FILE SCISEARCH
2	FILE TOXCENTER
4	FILE USPATFULL
2	FILE WPIDS
2	FILE WPINDEX

23 FILES HAVE ONE OR MORE ANSWERS, 75 FILES SEARCHED IN STNINDEX

L1 QUE (BACTERIOPHAGE OR PHAGE) (w) 77

=> file hits	-dgene	-genbank
COST IN U.S. DOLLARS		
FULL ESTIMATED COST		

SINCE FILE	TOTAL
ENTRY	SESSION
1.14	1.35

FILE 'CAPLUS' ENTERED AT 18:49:37 ON 07 NOV 2004

USE IS SUBJECT TO THE TERMS OF YOUR STM CUSTOMER AGREEMENT.
PLEASE SEE "HELP USAGETERMS" FOR DETAILS.

COPYRIGHT (C) 2004 AMERICAN CHEMICAL SOCIETY (ACS)

FILE 'BIOSIS' ENTERED AT 18:49:37 ON 07 NOV 2004
COPYRIGHT (C) 2004 The Thomson Corporation.

FILE 'EMBASE' ENTERED AT 18:49:37 ON 07 NOV 2004
COPYRIGHT (C) 2004 Elsevier Inc. All rights reserved.

FILE 'MEDLINE' ENTERED AT 18:49:37 ON 07 NOV 2004

FILE 'USPATFULL' ENTERED AT 18:49:37 ON 07 NOV 2004
CA INDEXING COPYRIGHT (C) 2004 AMERICAN CHEMICAL SOCIETY (ACS)

FILE 'IFIPAT' ENTERED AT 18:49:37 ON 07 NOV 2004
COPYRIGHT (C) 2004 IFI CLAIMS(R) Patent Services (IFI)

FILE 'SCISEARCH' ENTERED AT 18:49:37 ON 07 NOV 2004
Copyright (c) 2004 The Thomson Corporation.

FILE 'BIOTECHABS' ACCESS NOT AUTHORIZED

FILE 'BIOTECHDS' ENTERED AT 18:49:37 ON 07 NOV 2004
COPYRIGHT (C) 2004 THE THOMSON CORPORATION

FILE 'DISSABS' ENTERED AT 18:49:37 ON 07 NOV 2004
COPYRIGHT (C) 2004 ProQuest Information and Learning Company; All Rights Reserved.

FILE 'ESBIOSASF' ENTERED AT 18:49:37 ON 07 NOV 2004
COPYRIGHT (C) 2004 Elsevier Science B.V., Amsterdam. All rights reserved.

FILE 'LIFESCI' ENTERED AT 18:49:37 ON 07 NOV 2004
COPYRIGHT (C) 2004 Cambridge Scientific Abstracts (CSA)

FILE 'PASCAL' ENTERED AT 18:49:37 ON 07 NOV 2004
Any reproduction or dissemination in part or in full,
by means of any process and on any support whatsoever,
is prohibited without the prior written agreement of INIST-CNRS.
COPYRIGHT (C) 2004 INIST-CNRS. All rights reserved.

FILE 'PROMT' ENTERED AT 18:49:37 ON 07 NOV 2004
COPYRIGHT (C) 2004 Gale Group. All rights reserved.

FILE 'TOXCENTER' ENTERED AT 18:49:37 ON 07 NOV 2004
COPYRIGHT (C) 2004 ACS

FILE 'WPIDS' ENTERED AT 18:49:37 ON 07 NOV 2004
COPYRIGHT (C) 2004 THE THOMSON CORPORATION

FILE 'WPINDEX' ACCESS NOT AUTHORIZED

FILE 'ADISINSIGHT' ENTERED AT 18:49:37 ON 07 NOV 2004
COPYRIGHT (C) 2004 Adis Data Information BV

FILE 'BIOENG' ENTERED AT 18:49:37 ON 07 NOV 2004

COPYRIGHT (C) 2004 Cambridge Scientific Abstracts (CSA)

FILE 'BIOTECHNO' ENTERED AT 18:49:37 ON 07 NOV 2004
COPYRIGHT (C) 2004 Elsevier Science B.V., Amsterdam. All rights reserved.

FILE 'PHAR' ENTERED AT 18:49:37 ON 07 NOV 2004
COPYRIGHT (C) 2004 PJB Publications Ltd. (PJB)

=> s 11

L2 59 L1

=> dup rem l2
DUPLICATE IS NOT AVAILABLE IN 'ADISINSIGHT, PHAR'.
ANSWERS FROM THESE FILES WILL BE CONSIDERED UNIQUE
PROCESSING COMPLETED FOR L2
L3 28 DUP REM L2 (31 DUPLICATES REMOVED)

=> s 13 not pelletier?/au
NUMERIC VALUE NOT VALID 'PELLETIER?'
NUMERIC VALUE NOT VALID 'PELLETIER?'
L4 22 L3 NOT PELLETIER?/AU

=> s 14 and py<=1998
<-----User Break----->

SEARCH ENDED BY USER

=> s 14 and sequence
L5 5 L4 AND SEQUENCE

=> d 15 bib ab 1-5

L5 ANSWER 1 OF 5 CAPLUS COPYRIGHT 2004 ACS ON STM
AN 2003:6130 CAPLUS
DN 138:69476
TI Method for separating bacterial cells and bacterial cell components using
IMmobilized bacteriophage or bacteriophage proteins
IN Karelh, Michael Schuetz; Grassl, Renate; Meyer, Roman; Frick, Sibylle;
Robl, Ingrid; Zander, Thomas; Miller, Stefan
PA Profes A.-G., Germany
SO PCT Int. Appl., 40 pp.
CODEN: PIXXD2
DT Patent
LA German
FAN CNT 1

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
PI WO 2003000888	A2	20030103	WO 2002-DE2302	20020624
WO 2003000888	A3	20031224		
W:	AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GR, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MY, NZ, OM, PH, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, ST, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZM, ZW, AM, AZ, BY, KG, KZ, MD, RU, TJ, TM			
RM:	GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW, AT, BE, CH,			

CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, TR, BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG DE 10129815 A1 20030109 DE 2001-10129815 20010624 EP 1399551 A2 20040324 EP 2002-750798 20020624 R: AT, BE, CH, DE, DK, ES, FR, GB, GR, IT, LI, LU, NL, SE, MC, PT, IE, SI, LT, LV, FI, RO, MK, CY, AL, TR PRAI DE 2001-10129815 A 20010624 WO 2002-DE2302 W 20020624

AB The invention relates to a method for selectively sepg. bacterial cells and/or cell components such as plasmids, whereby the sepg. is carried out using a solid support to which bacteriophages or bacteriophage proteins are attached. Thus, biotin-labeled phage T4 were immobilized on streptavidin-coated magnetic beads and these were used in affinity purifn. of E. coli. Plasmids of the purified E. coli may then be isolated.

L5 ANSWER 2 OF 5 BIOTECHDS COPYRIGHT 2004 THE THOMSON CORP. ON STN AN 2003-28040 BIOTECHDS

TI Adapter-directed display system having expression vector with ***sequence*** encoding exogenous polypeptide and helper vector, for displaying exogenous polypeptide e.g., receptor ligand on outer surface of genetic package; recombinant protein production via plasmid expression in host cell

AU WANG C; ZHONG P; WANG X

PA WANG C; ZHONG P; WANG X

PI US 200104355 5 Jun 2003

AI US 2001-33399 2 Nov 2001

PRAI US 2001-33399 2 Nov 2001; US 2001-33399 2 Nov 2001

DT Patent

LA English

OS WPI: 2003-801222 (75)

AB DERIVAT ABSTRACT.

NOVELTY - An adapter-directed display system for displaying an exogenous polypeptide on the outer surface of a genetic package, comprising an expression vector with a ***sequence*** that encodes exogenous polypeptide fused to first adapter ***sequence***, and a helper vector with outer-surface sequences encoding proteins fused to second adapter, the polypeptide is produced in a host cell to cause display of the polypeptide, is new.

DETAILED DESCRIPTION - An adapter-directed display system (1) for displaying an exogenous polypeptide on the outer surface of a genetic package, comprising an expression vector comprising a coding ***sequence*** that encodes the exogenous polypeptide fused in-frame

to a first adapter ***sequence***, where the vector is devoid of outer-surface sequences encoding any functional outer-surface proteins of the genetic package, and a helper vector comprising outer-surface sequences encoding outer-surface proteins necessary for packaging the genetic package, where at least one of the outer-surface protein is fused in-frame to a second adapter, the first and second adapter acting, when the polypeptide is produced in a suitable host cell, to cause the display of the polypeptide through pairwise interaction between the first and second adapters. INDEPENDENT CLAIMS are also included for the following:

(1) a helper vector (II) for displaying a polypeptide on the outer surface of a genetic package comprising, outer-surface sequences necessary for packaging the genetic package, where at least one of the surface presenting sequences in fused in-frame to an adapter, the adapter acting, when the polypeptide is produced in a suitable host cell, to

cause the display of the polypeptide; (2) an expression vector (III) for producing a polypeptide within or on the outer surface of a genetic package, comprising a coding ***sequence*** encoding the polypeptide fused in-frame to a first adapter, where the vector is devoid of outer-surface sequences encoding any functional outer-surface proteins of the genetic package, and expression of the polypeptide on the outer surface of the genetic package is mediated through non-covalent pairwise interaction between the first adapter and a second adapter, where the second adapter is fused to an outer-surface protein; (3) a kit comprising (I), (II) or (III) in suitable packaging; (4) a host cell comprising (I), (II) or (III); (5) a polypeptide displayed on the outer surface of a genetic package using (I); (6) a genetic package (IV) displaying on its outer surface a fusion polypeptide, the fusion polypeptide comprising a polypeptide ***sequence*** to be displayed, fused in-frame with a first adapter, the first adapter acting, when the fusion polypeptide is produced in a suitable host cell, to cause the display of the fusion polypeptide through non-covalent pairwise interaction between the first adapter and a second adapter that is linked to an outer-surface protein; (7) a selectable library comprising several genetic packages, at least one being (IV); (8) a selectable library (V) comprising several genetic packages, at least one member of several packages displaying a polypeptide on its outer surface using (I); and (9) detecting (VI) the presence of a specific interaction between a test agent and an exogenous polypeptide that is displayed on a genetic package, involves providing a genetic package displaying the exogenous polypeptide that is prepared using (I), contacting the genetic package with the test agent under conditions suitable to produce a stable polypeptide-agent complex, and detecting the formation of the stable polypeptide-agent complex on the genetic package, thus detecting the presence of a specific interaction.

BIOTECHNOLOGY - Preferred System: (1) is a phage or bacterial display system. The genetic package is chosen from viruses, cells and spores. The outer-surface sequences encode functional coat proteins of a phage. The phage is a filamentous phage. The outer-surface sequences are chosen from gene III, gene VI, gene VII, gene VIII and gene IX of a filamentous phage. The outer-surface sequences encode bacterial outer-surface proteins chosen from Lpp-OmpA, Trar, Pal, OprI, Imp and AIDA-1. The first and second adapters are homodimerization or heterodimerization sequences. The homodimerization sequences consist of a pair of cysteine residues. The first and second adapters form a coiled-coil dimer, where the first and second adapters are leucine zippers and comprise heterodimeric receptor sequences that mediate heterodimerization of the receptors. The first and second adapters comprise heterodimerization sequences of GABAR receptor 1/receptor 2 and GABAR receptor 2/receptor 1, respectively. The helper vector further comprises at least one additional copy of outer-surface ***sequence*** that competes for packaging with the fusion outer-surface ***sequence***.

sequence. The expression vector is chosen from pBAMX14 and pBAMX15. The phage helper vector is chosen from GM-Ultrahelper phage vector, CM-Ultrahelper phage vector and GMCT-Ultrahelper phage vector. Preferred Vector: (II) is a phage helper vector or bacterial helper vector. The adapter causes the display of the polypeptide through pairwise interaction with a second adapter, which is fused in-frame with the polypeptide. The adapter causes the display of the polypeptide in the absence of expression of an outer-surface protein through a phagemid vector or a plasmid. The two adapters are heterodimerization or homodimerization sequences. (III) is a phagemid vector or bacterial expression vector. The outer-surface sequences encode bacterial

outer-surface proteins. Preferred Method: In (M1), the exogenous polypeptide is chosen from antigen-binding unit, cell surface receptor, receptor ligand, cytosolic protein, secreted protein, and nuclear protein. Preferably, the exogenous polypeptide is antigen-binding unit. The test agent is chosen from protein, polysaccharides, lipid and their combinations.

USE - (1) is useful for displaying a polypeptide on the outer surface of a genetic package, which involves causing (1) to be transcribed and translated in a suitable host cell. (V) is useful for obtaining a polypeptide with desired property, which involves providing (V), and screening the selectable library to obtain at least one genetic package displaying a polypeptide with the desired property. The desired property is binding specificity to an agent of interest. Screening the selectable library further involves isolating the genetic package that displays a polypeptide having the desired property. Isolating the genetic package further involves obtaining a nucleotide ***sequence*** from the genetic package that encodes the polypeptide with the desired property. The polypeptide with the desired property is chosen from antigen-binding unit, cell surface receptor, receptor ligand, cytosolic protein, secreted protein, nuclear protein, and their functional motif (claimed).

EXAMPLE - Construction of an adapter-directed display system comprising CM-Ultrahelper phage vector was done as follows. The PABM13 vector was constructed by replacing the ***sequence*** between the XbaI and BglII sites of vector PABMD1 with a synthetic DNA fragment comprising 5'-3' gene III leader ***sequence***, a KpnI site, a coding ***sequence*** for Ala-Cys-Gly-Gly and a Myc-tag. The synthetic ***sequence*** was linked in-frame with gene III in PABMD1 vector. The CM-Ultrahelper phage vector was constructed by replacing the KpnI/BamHI fragment encoding and partial p III protein in the KOT Kpn helper vector with the corresponding fragment encoding a partial p III leader and the adapter2-pIII fusion protein from the PABM13 vector. The resulting CM-Ultrahelper phage vector encoded an engineered pIII capsid fused with Cys-Myc domain placed at the N-terminal of pIII. The B9 KOT kpn helper phage clone was used for the construction of CM-Ultrahelper phage vector. After subcloning of the engineered gene III fragment into KOT kpn phage vector, 24 kanamycin-resistant colonies were grown in 96-well microtiter plates in 2xYT medium with 70 microg/ml kanamycin, and supernatants were used for phage screening by phage enzyme linked immunosorbent assay (ELISA). Twenty three clones were found to generate phage particles. To confirm the ability of CM-Ultrahelper phages to package the expressed Cys-Myc-pIII fusion into the phage particles, ELISA assay using anti-Myc antibody was performed. The CM-Ultrahelper phages from the selected five clones displayed the Myc-tag on their surfaces. Such Myc-tag was not detectable in the KOT helper phage negative control. The vector PABM15 and was constructed from PABMD2 by replacing the fd gene III fragment of PABMD2 with a synthetic DNA fragment encoding the HA-Tag and Gly-Gly-Cys. To demonstrate that a functional protein was displayed using CM-Ultrahelper phage vector, the single-chain antibody AM1 was subcloned into the PABM15 vector. The resulting PABM15-AM1 was transformed into TG1 cells, and the cells were superinfected with CM-Ultrahelper phage with a multiplicity of infection (MOI) of 100. Phage particles were generated and purified. The single-chain antibody displayed on the phage surface was detected by phage ELISA using plates coated with AM1-antigen 2x1012 phages were added for each well. The secondary antibody was horse radish peroxidase (HRP)-conjugated anti-M13 antibody. The ELISA results showed that the phagemid particles generated

from all four MOI infections were capable of specifically binding to AM1-antigen, which indicated that functional single-chain antibody was displayed on the phage surface. The control phagemids generated from TG1 carrying PABM15-AM1/M13K07 vectors did not bind to AM1-antigen. The phage particles were also used for western blot analysis. Phage particles were denatured by heating in sodium dodecyl sulfate (SDS) sample buffer under non-reducing conditions, the scfv antibody was only displayed by CM-Ultrahelper phage, not by M13K07 helper ***phage***. (****7*** pages)

L5 ANSWER 3 OF 5 DISSABS COPYRIGHT (C) 2004 ProQuest Information and Learning Company: All Rights Reserved on STN
AN 84:22121 DISSABS Order Number: AAR8501756
TI THE ISOLATION AND CHARACTERIZATION OF THE BACILLUS SUBTILIS GENE ENCODING
THE MAJOR SIGMA FACTOR OF RNA POLYMERASE (SIGMA-43, SIGMA-55, RPOD)
AU GITT, MICHAEL ALAN [Ph.D.]
CS UNIVERSITY OF CALIFORNIA, DAVIS (0029)
SO Dissertation Abstracts International, (1984) Vol. 45, No. 11B, p. 3493.
Order No.: AAR8501756. 156 pages.

DT Dissertation
FS DAI
LA English
ED Entered STN: 19921118
AB Last Updated on STN: 19921118

To understand the in vivo role of the major sigma subunit of Bacillus subtilis RNA polymerase, and to understand more about its mechanism of action and regulation, we have isolated its gene, the rpoD gene. Using immunological screening, we isolated 13 phage, all having at least two 3.5 kb EcoRI fragments in common. One representative phage, (lambda)GWS-sigma-82, contained only the two 3.5 kb inserts in addition to the lambda arms. Its lysate was shown to contain a product which crosses-reacted with anti-sigma-43 (aka sigma-55) antibody, and which was identical to authentic sigma-43 in apparent molecular weight, charge, and VS protease peptide map.

Restriction fragments of the insert DNA were subcloned into plasmid pCP12 to localize the rpoD gene to a 1.7 kb EcoRI-SphI fragment, using a maxicell assay. Other proteins were encoded in the adjacent DNA within the original 7 kb insert. The subcloned 1.7 kb fragment coded for a product which was identical to authentic sigma-43 in apparent molecular weight, peptide map, and immunologic properties.

The 1.7 kb fragment was sequenced and the primary structure of sigma-43 deduced and compared to the N-terminal ***sequence*** of authentic sigma-43. The molecular weight of sigma-43 is calculated as 42,828 daltons, rather than the 55,000 value obtained from gel electrophoresis. I have located a strong ribosomal binding site just upstream of the initiator methionine and a rho-independent transcription terminator ***sequence*** just downstream of the translational termination codon. Codon usage resembles the average for several B. subtilis genes. The rpoD gene appears to be part of an operon, as is the case in E. coli.

The sigma-43 protein exhibits extensive homology, especially in the C-terminal two-thirds of the ***sequence***, to sigma-70 of E. coli. A large internal stretch of 245 amino acids present in sigma-70 is absent in sigma-43. Other proteins exhibiting homology with sigma-43 include the core-binding DNA protein, the heat shock regulator protein of E. coli, and specific RNA-binding proteins. No amino acid homology was found between sigma-43 and B. subtilis phage SP01 sigma-gp28, E. coli

phage **77*** RNA polymerase, or E. coli DNA primase.

L5 ANSWER 4 OF 5 PROMT COPYRIGHT 2004 Gale Group on STN

AN 2003:538870 PROMT
 TI Maximizing the yield of full-length RNA from an in vitro transcription
 reaction. (Tools & Techniques)
 AU Mele, Ron; Pease, Jim; Kramer, Katharine
 SO Biotechnology, (Sept 2003) Vol. 28, No. 9, pp. 8(2).
 PB Reed Business Information
 DT Newsletter
 LA English
 WC 1986

AB *FULL TEXT IS AVAILABLE IN THE ALL FORMAT*
 Abstract
 THIS IS THE FULL TEXT: COPYRIGHT 2003 Reed Business Information
 Subscription: \$37.90 per year. Published monthly. Box 6399, Torrence, CA
 90504.

L5 ANSWER 5 OF 5 WPIDS COPYRIGHT 2004 THE THOMSON CORP ON STN
 AN 1987-286263 (41) WPIDS
 CR 1993-257894 (32)
 DMC C1987-121376
 TI Pure nucleic acid comprising G 10 L ***sequence*** - useful for
 enhancing expression of genes in bacteria for prodn. of hormones, plant or
 bacterial enzymes etc..

DC B04 D16
 IN OLINS, P O; OLAFS, P
 PA (MONS) MONSANTO CO
 CYC 21

PI EP 24146 A 19871014 (198741)* EN 32
 R: AT BE CH DE ES FR GB GR IT LI LU NL SE
 AU 8770661 A 19871001 (198746)
 HU 43625 T 19871130 (198751)
 JP 63000289 A 19880105 (198806)
 DK 8701544 A 19871221 (198810)
 ZA 8702242 A 19871124 (198817)
 EP 24146 B1 19950301 (199513) EN 39
 R: AT BE CH DE ES FR GB GR IT LI LU NL SE
 DE 3751100 G 19950406 (199519)
 ES 2070828 T3 19950616 (199531)
 IL 82017 A 19960119 (199616)
 IE 67890 B 19960501 (199629)
 JP 2511948 B2 19960703 (199631) 32
 CA 1340091 C 19981020 (199901)

ADT EP 24146 A EP 1987-870039 19870326; JP 63000289 A JP 1987-72974 19870326;
 ZA 8702242 A ZA 1987-2242 19870326; EP 24146 B1 EP 1987-870039 19870326;
 DE 3751100 G DE 1987-3751100 19870326; IL 82017 A IL 1987-82017 19870326; ES 2070828
 T3 EP 1987-870039 19870326; JP 2511948 B2 JP 1987-72974 19870326; CA 1340091 C
 IE 1987-782 19870326; JP 2511948 B2 JP 1987-72974 19870326; CA 1340091 C
 DE 3751100 G Based on EP 24146; ES 2070828 T3 Based on EP 24146; JP
 FDT 2511948 B2 Previous Publ. JP 63000289
 PRAI US 1987-5821 19870204; US 1986-845159 19860327
 AB EP 24146 A UPAB: 19970502

(1) Pure nucleic acid molecule comprising a G10L ***sequence*** is

new. The ***sequence*** esp. comprises about the first 100 nucleotides
 immediately 5' to the translation start codon of the ***bacteriophage***
 ****77*** gene 10 coding ***sequence***. (2) Prodn. of a
 heterologous
 protein (1) in bacteria comprises causing expression of a gene comprising
 a G10L ***sequence*** operatively joined to a DNA ***sequence***
 coding for (1).
 USE/ADVANTAGE - The nucleic acid molecule enhances the expression of
 a wide variety of genes, both prokaryotic and eucaryotic, in bacteria. (1)
 is esp. a mammalian growth hormone, atriopeptigen or plant or bacterial
 enzyme.
 Dwg. 0/12

>> d his

(FILE 'HOME' ENTERED AT 18:48:02 ON 07 NOV 2004)
 INDEX 'ADISCTI, ADISINSIGHT, ADISNEWS, AGRICOLA, ANABSTR, ANTE, AQUALINE,
 AQUASCI, BIOBUSINESS, BIOCOMMERCE, BIOENG, BIOSIS, BIOTECHABS, BIOTECHDS,
 BIOTECHNO, CAB, CANCERLIT, CAPLUS, CEABA-VTB, CIN, CONFSCI, CROPB,
 CROPU, DDFB, DGENE, DISSABS, ...' ENTERED AT 18:48:16 ON 07 NOV 2004
 SEA (BACTERIOPHAGE OR PHAGE) (W) 77

1 FILE ADISINSIGHT
 1 FILE BIOENG
 10 FILE BIOSIS
 2 FILE BIOTECHABS
 2 FILE BIOTECHDS
 1 FILE BIOTECHNO
 1 FILE CAPLUS
 25 FILE DGENE
 2 FILE DISSABS
 4 FILE EMBASE
 2 FILE ESBIOBASE
 24 FILE GENBANK
 3 FILE IFIPAT
 2 FILE LIFESCI
 4 FILE MEDLINE
 1 FILE PASCAL
 2 FILE PHAR
 2 FILE PROMT
 3 FILE SCISEARCH
 2 FILE TOXCENTER
 4 FILE USPATFUL
 2 FILE WPIDS
 2 FILE WPINDEX
 2 QUE (BACTERIOPHAGE OR PHAGE) (W) 77

FILE 'CAPLUS, BIOSIS, EMBASE, MEDLINE, USPATFUL, IFIPAT, SCISEARCH,
 BIOTECHDS, DISSABS, ESBIOBASE, LIFESCI, PASCAL, PROMT, TOXCENTER, WPIDS,
 ADISINSIGHT, BIOENG, BIOTECHNO, PHAR' ENTERED AT 18:49:37 ON 07 NOV 2004
 59 S L1
 28 DUP REM L2 (31 DUPLICATES REMOVED)
 22 S L3 NOT PELLETIERER/AU
 5 S L4 AND SEQUENCE

```

=> log h
COST IN U.S. DOLLARS
FULL ESTIMATED COST
DISCOUNT AMOUNTS (FOR QUALIFYING ACCOUNTS)
CA SUBSCRIBER PRICE

```

	SINCE FILE ENTRY	TOTAL SESSION
	49.63	50.98
	-0.70	-0.70

SESSION WILL BE HELD FOR 60 MINUTES
 STN INTERNATIONAL SESSION SUSPENDED AT 18:51:54 ON 07 NOV 2004